What is claimed is:

5

10

15

20

- 1. An isolated protein comprising an amino acid sequence as set forth in SEQ ID NO: 2, 4, 6 or 8.
- 2. An isolated protein consisting of an amino acid sequence as set forth in SEQ ID NO: 2, 4, 6 or 8.
 - 3. An isolated protein, wherein the protein comprises a DNA repair activity and is encoded by a nucleic acid that hybridizes under stringent conditions with a nucleic acid comprising all or a part of the nucleotide sequence as set forth in SEQ ID NO: 1, 3, 5 or 7.
 - 4. An isolated nucleic acid encoding a protein comprising an amino acid sequence as set forth in SEQ ID NO: 2, 4, 6 or 8.
 - 5. An isolated nucleic acid comprising: (a) a DNA consisting of the nucleotide sequence as set forth in SEQ ID NO: 1, 3, 5 or 7; or, a complementary strand to (a).
 - 6. An isolated nucleic acid, wherein the nucleic acid encodes a polypeptide comprising a DNA repair activity and hybridizes under stringent conditions with a nucleic acid comprising the nucleotide sequence as set forth in SEQ ID NO: 1, 3, 5 or 7, or, with a complementary strand thereto.
 - 7. An isolated nucleic acid, wherein the nucleic acid encodes a polypeptide comprising a DNA repair activity and hybridizes under stringent conditions with a probe prepared from a nucleic acid comprising all or a part of a nucleotide sequence as set forth in SEQ ID NO: 1, 3, 5 or 7, or from a complementary strand thereto.
 - 8. The isolated nucleic acid, wherein the stringent conditions comprise a wash step comprising a wash in 0.2X SSC at a temperature of about 65°C for about 15 minutes.

- 9. A recombinant vector comprising a nucleic acid as set forth in claims 4, 5, 6 or 7.
- 10. A recombinant vector comprising a nucleic acid encoding a polypeptide as set forth in claims 1, 2 or 3.

10

15

20

25

30

- 11. A transformed cell comprising a recombinant vector as set forth in claim 9.
- 12. A transformed cell comprising the recombinant vector as set forth in claim 10.
- 13. A method of producing a DNA repair enzyme, comprising
- (a) culturing a transformed cell according to claim 11 or claim 12, and
- (b) recovering the DNA repair enzyme from the resultant culture, thereby producing a DNA repair enzyme.
 - 14. A method of producing a DNA repair enzyme, comprising
 - (a) culturing a transformed cell according to claim 11 or claim 12, and
- (b) recovering the DNA repair enzyme from the resultant culture, thereby producing a DNA repair enzyme.
- 15. A method of repairing a DNA for sequence errors or base mismatch errors, comprising carrying out a DNA synthesis reaction in the presence of a polypeptide as set forth in claims 1, 2 or 3.
- 16. A method of preventing erroneous synthesis of DNA sequences, comprising carrying out a DNA synthesis reaction in the presence of a polypeptide as set forth in claims 1, 2 or 3.
- 17. A DNA repair gene-disrupted cell obtained by transferring into a host cell a nucleic acid as set forth in claims 4, 5, 6 or 7.

- 18. The DNA repair gene-disrupted cell of claim 17, wherein a modification gene has been incorporated into the nucleic acid.
- 19. The DNA repair gene-disrupted cell of claim 18, wherein the modification gene comprises a marker gene.
- 20. The DNA repair gene-disrupted cell of claim 17, wherein the host is a bacterium.
- 21. The DNA repair gene-disrupted cell of claim 20, wherein the bacterium is a thermophilic bacterium.
 - 22. The DNA repair gene-disrupted cell of claim 21, wherein the thermophilic bacterium is a bacterium of the genus *Thermus*.
 - 23. The DNA repair gene-disrupted cell of claim 22, wherein the thermophilic bacterium is a *Thermus thermophilus*.
 - 24. An array comprising a nucleic acid as set forth in SEQ ID NO: 1, 3, 5 or 7.
 - 25. An array comprising a nucleic acid as set forth in claims 4, 5, 6 or 7.
 - 26. A method of screening a composition for its ability to specifically bind to a DNA repair enzyme comprising:
 - (a) contacting the a DNA repair enzyme with the composition, wherein the DNA repair enzyme is a polypeptide encoded by a nucleic acid sequence as set forth in claims 4, 5, 6 or 7; and,
 - (b) determining if the composition specifically binds to the DNA repair enzyme.

10

. :

15

20

- 27. A method for inhibiting the expression of a DNA repair enzyme encoding nucleic acid in a cell, the method comprising the following steps:
- (a) providing a nucleic acid operably linked to a promoter that expresses an inhibitory sequence, wherein the inhibitory sequence comprises all or part of a nucleic acid sequence as set forth in claims 4, 5, 6 or 7 and is expressed in a form sufficient to inhibit expression of a DNA repair enzyme message message; and,
- (b) expressing the inhibitory nucleic acid in an amount sufficient to inhibit the expression of the DNA repair enzyme encoding nucleic acid in the cell.
- 28. The method of claim 27, wherein the inhibitory sequence comprises an antisense sequence.
- 29. The method of claim 27, wherein the inhibitory sequence comprises a ribozyme sequence.
- 30. A method of expressing a heterologous nucleic acid sequence in a cell comprising:
- a) transforming the cell with a heterologous nucleic acid operably linked to a promoter, wherein the heterologous nucleic acid comprises a nucleic acid sequence as set forth in claims 4, 5, 6 or 7; and,
- b) growing the cell under conditions where the heterologous nucleic acid sequence is expressed in the cell.
- 31. A method for detecting a nucleic acid in a nucleic acid -containing biological sample, the method comprising the following steps:
- (a) contacting the sample with a nucleic acid probe comprising a nucleic acid sequence as set forth in claims 4, 5, 6 or 7;
 - (b) hybridizing the nucleic acid probe to the nucleic acid in the sample; and,
 - (c) detecting hybridization of the nucleic acids.

5

10

15

- 32. An fusion protein comprising a first amino acid sequence as set forth in SEQ ID NO: 2, 4, 6 or 8, or a subsequence thereof, and a second heterologous sequence.
- 33. An isolated antibody specifically reactive with a polypeptide as set forth in claim 1, claim 2 or claim 3 or a polypeptide encoded by a nucleic acid as set forth in claim 4, claim 5, claim 6, or claim 7.
 - 34. The antibody of claim 33, wherein the antibody is a monoclonal antibody.
- 10 35. A hybridoma cell comprising the monoclonal antibody of claim 34.